

Impact du jeûne du mois de Ramadan sur le profil lipidique et les facteurs de risque cardiovasculaire chez les coronariens stables.

Impact of Ramadan fasting on lipid profile and cardiovascular risk factors in patients with stable coronary artery disease

Habib Ben Ahmed¹, Emna Allouche¹, Kahena Bouzid², Sabine Zrelli², Wided Hmaid², Yosra molahedh², Wejdane Ouechtati¹, Leila Bezdah¹.

1 : Service de Cardiologie, Hôpital Charles Nicolle, Tunis, Tunisie.

2 : Service de Biochimie clinique, Hôpital Charles Nicolle, Tunis, Tunisie.

Résumé

Objectifs : l'objectif de cette étude était d'évaluer l'impact du Jeûne du Ramadan (JR) sur le profil lipidique et les facteurs de risque cardiovasculaire chez les patients atteints de coronaropathie stable.

Méthode : il s'agit d'une étude prospective portant sur des patients présentant une cardiopathie ischémique stable et ayant l'intention de jeûner le Ramadan, recrutés dans le service de cardiologie, hôpital Charles Nicolle, Tunisie durant le mois de ramadan de l'année 2020. Une évaluation des paramètres cliniques, des profils glycémiques et lipidiques, des taux de protéine C-réactive ultra-sensible et d'homocystéine a été effectuée avant et après le mois sacré.

Résultats : Cette étude n'a pas mis en évidence une altération significative du statut angineux. Les niveaux de cholestérol, de triglycérides, de lipoprotéine de faible densité et d'apolipoprotéine A ont été significativement améliorés après le JR par rapport à leurs valeurs de base. Il existe par ailleurs, une diminution significative de la glycémie à jeun, du taux d'insuline et de l'indice de résistance à l'insuline avec une baisse de la CRP us.

Conclusion : Chez les patients présentant une cardiopathie ischémique stable, le JR peut s'accompagner d'une amélioration du profil lipidique et des paramètres glycémiques et ne constitue pas un facteur d'augmentation des événements coronariens aigus.

Mots-clés

Facteurs de risque cardiovasculaire, jeûne, habitudes alimentaires, cardiopathie ischémique

Summary

Background: The effects of Ramadan fasting (RF) on cardiometabolic risk factors in patients with stable ischemic heart disease are not well known.

Aim : To evaluate the impact of RF on lipid profile and cardiovascular risk factors in patients with a stable coronary heart disease.

Methods: A prospective observational study carried out in the Cardiology department of Charles Nicolle Hospital (Tunisia). Eighty-four patients with a stable ischemic heart disease who intended to fast were enrolled during May 2020. Detailed clinical and biochemical assessments were performed before and after the holy month. Parameters of glycemic control, lipid profile, ultrasensitive C-reactive protein concentration (us-CRP) and homocysteine were performed before- and after- Ramadan (BR and AR, respectively).

Results: Eighty-four patients including 79 males and 5 females, with a mean age of 57±7 years completed the study. Levels of cholesterol, triglycerides, low-density lipoprotein-cholesterol and apoprotein A were significantly improved AR fasting in comparison with their BR values. There was a significant decrease in blood fasting glucose, insulin level, Homeostasis model assessment of insulin resistance index and in us-CRP level.

Conclusion: In patients with stable ischemic heart disease, RF may be accompanied by an improvement of lipid profile and glycemic parameters without increase in coronary events.

Keywords

Cardiovascular Risk factors, fasting, eating habits, ischemic heart disease.

Correspondance

Habib Ben Ahmed

Department of cardiology, Charles Nicolle Hospital, Tunis, Tunisia

eMail: dr.benahmed.habib@gmail.com

INTRODUCTION

Ramadan fasting (RF) concerns over one billion observant Muslims in the world [1]. From a physiological standpoint, Islamic fasting is a unique fasting model. It is distinct from voluntary or experimental fasting. During Ramadan, fasting Muslims carefully abstain from all oral intakes of food, water, beverages, and drugs from sunrise to sunset each day during a whole month.

RF traditionally makes a radical change in lifestyle and dietary habits which may affect cardiovascular risk in patients with coronary artery disease (CAD) [2]. Although more than a billion Muslims worldwide fast during Ramadan, there is no overwhelming consensus on its effects on ischemic heart disease [3]. Some published studies have shown that the effects of fasting on stable patients with CAD are minimal and more than 70% of them can experience RF with no clinical deterioration [4]. In fact, patients with cardiovascular disease are categorized as high risk for RF in the last practical guidelines [5]. However, several patients insist to fast although doctor's advice.

Otherwise, most of these studies were performed in the Middle East and Gulf areas, and therefore such conclusions may not be extrapolated to the Mediterranean region's population, where the period of daily fasting may be longer and the quality of food intake and physical activity may change [2,6-8]. In the present study, the aim of this study was to evaluate the potential impact of fasting on lipid profile and other biochemical markers that are traditionally used for accurately predicting cardiovascular events in patients with a stable coronary artery disease.

METHODS

Study design

A prospective observational study carried out in the Cardiology department of Charles Nicolle Hospital. Eighty-four outpatients with a stable ischemic heart disease with the deliberate intention to fast were enrolled. Detailed clinical and biochemical assessments were performed a month before the beginning of Ramadan and within the first week after this month.

This study was conducted in the month of Ramadan during May 2020 in the city of Tunis, Tunisia. Patients were followed with three consecutive visits BR, during the first week of Ramadan to witness whether they were capable to continue fasting, and immediately after the sacred month.

The average fasting period totals was 16.5hours. Height and weight were recorded for each patient and averaged for analysis. Body mass index was calculated to the nearest decimal place by using the formula weight in kilograms/ height in square meters. Blood pressure was measured by auscultatory method in the seated position, with a cuff bladder adapted to the patient's arm circumference.

Patients

The study enrolled patients between 18 and 70 years old with a documented previous history of coronary artery disease (CAD) in the past 5 years. Patients with ongoing myocardial ischemia (class III-IV angina of The *Canadian Cardiovascular Society*) [9], pregnant and lactating women, and patients aged below 18 years old and those who fasted for less than 20 days were excluded. Patients with any type of acute or chronic inflammation were also excluded. Written informed consent was obtained from all patients. All patients were instructed not to make any changes to their usual physical activities during Ramadan. Ethical committee approval for this study was obtained.

Sample collection and preservation

Fasting blood samples were collected from each patient between 9 am and 11 am, a week before Ramadan and in the first 7 days after the end of this month. Approximately 6 ml venous blood sample was collected from the left arm by using a vacuum sampling method and analyzed in the same hospital affiliated laboratory. Blood samples were centrifuged to separate the plasma at 3000 rpm for 5 min at 4 C and were kept at -20 C until analysis.

Determination of biochemical parameters

The levels of triglycerides, total cholesterol, high density lipoprotein-cholesterol (HDL-CT), apoproteins A and B, and total protein were measured by an enzymatic colorimetric method with an automated chemical analyzer (The Abbott Architect c8000).

Low density lipoprotein-cholesterol (LDL-CT) was estimated following the Friedewald formula, HDL risk was also estimated [Cholesterol (mmol /l) /HDL-CT (mmol/l)] [10].

The level of fasting glycaemia was measured using the glucose oxidase method and noted as mmol/l. Insulin concentration was measured by chemiluminescence.

Homeostasis model assessment of insulin resistance index (HOMA-IR) was calculated using fasting glucose and fasting insulin as follows: [fasting insulin (mUI/L) fasting glucose (mmol/l)] / 22.5. HOMA-IR \geq 2.6 was considered a marker of insulin resistance[11]. Ultrasensitive C-reactive protein (Us-CRP) and Homocysteine were measured using a radioimmunoassay method (automate).

Statistical analyses

The data were reported as mean \pm standard deviation. A paired t-test was used to compare pre- and post-Ramadan levels of biochemical variables. Statistical analysis was performed using SPSS 20.0 software. Differences were considered significant at $p < 0.05$.

RESULTS

A total of 114 patients initially consented to voluntarily participate in the study; however, 22 patients were excluded because of fasting for less than 20 days and 8 patients declined to participate for personal reasons. Eighty-four patients including 79 males and 5 females with a mean age of 57 ± 7 years completed the study. During the holly month we did not identify any clinical deterioration or significant worsening of biochemical parameters. Baseline clinical characteristics, history and ongoing treatment are summarized in Table 1.

The values of total cholesterol, triglycerides, and LDL-CT were significantly decreased after RF; however, the changes in HDL-CT cholesterol were not significant. HDL risk decreased significantly during Ramadan than before fasting (4.85 ± 1.49 vs 4.26 ± 1.37 $p < 0.001$). Apoprotein A significantly increased after fasting, but we didn't find any significant changes in apoprotein B before and after fasting (Table 2).

Fasting glycaemia, insulin concentrations and HOMA-IR index showed a significant decrease after RF ($p < 0.05$) (Fig. 1). Fasting decreased significantly glycaemia by 8%. After-RF, insulin levels were lower than before-RF levels (11.69 ± 6 vs 9.93 ± 5.13 $\mu\text{U/ml}$, $p < 0.001$).

Also, the HOMA-IR index significantly decreased after Ramadan (3,84 vs 3,08 %, $p < 0,001$).

Table1: Baseline clinical characteristics, medical history and treatment

Cardiovascular risk factors	84 patients ,nb(%)
Hypertension	57(68%)
Diabetes	26(31%)
Dyslipidemia	18(13%)
Smoking	60(71%)
Family coronary artery disease	20(24%)
Sedentary lifestyle	62(74%)
Cardiovascular history	
Stroke	5(6%)
Myocardial infarction	18(21%)
Coronary angioplasty	26(31%)
Coronary artery bypass graft	27(32%)
Treatment	
Statins	47 (56%)
Oral antidiabetics	26(31%)
Angiotensin Converting enzyme inhibitor	52 (62%)
Beta-blocker	57(68%)
Calcium inhibitor	7 (8%)
Diuretics	5(6%)
Aspirin	72 (86%)
Clopidogrel	11(13%)

Table 2: Effects of Ramadan fasting on clinical ,anthropometric and biochemical parameters

Variables	Pre-Ramadan	Post-Ramadan	p value	Normal values
BMI (Kg/m ²)	26.84 \pm 3.18	26.80 \pm 3.14	0.094	18.5-25
SBP (mmHg)	132.9 \pm 16	129.9 \pm 17	0.03	
DBP (mmHg)	83 \pm 8	82 \pm 7	0.07	
Cholesterol (mmol/l)	4.34 \pm 1.2	3.83 \pm 1.18	<0.001	3.2-5.18
Triglycerides (mmol/l)	2.04 \pm 1.34	1.79 \pm 1.33	<0.05	0.4-1.7
LDL-CT (mmol/l)	2.51 \pm 0.91	2.26 \pm 0.87	<0.001	<4.1
HDL-CT (mmol/l)	0.91 \pm 0.18	0.91 \pm 0.16	0.750	>1.3
HDL- risk	4.85 \pm 1.49	4.26 \pm 1.37	<0.001	
Apoprotein A (g /l)	1.21 \pm 0.15	1.26 \pm 0.15	<0.01	1.11-1.65
Apoproteins B (g /l)	0.85 \pm 0.27	0.97 \pm 0.53	0.055	0.69-1.09
Fasting Glucose (mmol/l)	7.14 \pm 3.21	6.58 \pm 2.68	<0.001	3.89-5.83
Insulin(μ U/ml)	11.69 \pm 6	9.93 \pm 5.13	<0.001	10-20
HOMA-IR(%)	3.84 \pm 2.8	3.08 \pm 2.5	<0.001	
Homocysteine (μ mol/l)	11.18 \pm 1.79	11.4 \pm 1.68	0.159	4.6-12.44
us-CRP (mg/l)	2.18 \pm 1.22	1.97 \pm 1.08	0.034	<8

Data was represented as mean \pm SD. SD: standard deviation, BMI: body mass index, SBP: systolic blood pressure, DBP: diastolic blood pressure. HDL-CT: High density lipoprotein-cholesterol, LDL-CT: Low density lipoprotein-cholesterol ,us-CRP :Ultra-sensitive C-reactive, protein HOMA-IR index :Homeostasis model assessment of insulin resistance index.

Ultra-sensitive CRP decreased significantly during Ramadan ($p=0.034$) (Fig. 1) but homocysteine didn't significantly change after fasting (11.18 ± 1.79 vs 11.40 ± 1.68 ; $p = 0.159$).

DISCUSSION

This study demonstrates that RF can lead to an improvement of lipid profile and glycemic parameters and does not increase acute coronary events in patients with stable ischemic heart disease.

Muslims worldwide represent a diverse and heterogeneous population in terms of health status. Many Muslim patients insist on fasting Ramadan Although Islam exempts them from fasting, especially when it could worsen one's illness or delay recovery. Major changes in meal timings during fasting days, as well as changes in the quality and quantity of diet, could affect metabolic profiles.

Our results showed that fasting effects are beneficial and the majority of patients with stable coronary artery disease can endure RF without any clinical deterioration or any significant worsening of their anginal status.

These results are consistent with the investigation of Chamsi-Pasha et al which reported a non-significant effect of fasting on hospitalization rates for heart failure, acute myocardial infarction, and the incidence of angina pain in patients with stable cardiovascular diseases[12]. Also, Khafaji et al reported no adverse effects on the clinical status of stable cardiac patients while fasting during Ramadan [13]. Another study showed that there was no significant difference in the number of hospitalizations for heart failure while fasting in Ramadan when compared to the non-fasting months[4].

Similar to our results, Mossavi and al showed that patients with underlying stable coronary heart disease can safely observe fasting in Ramadan without an increased incidence of acute coronary syndromes[14].

The present study showed a significant decrease in cholesterol, triglycerides, LDLc, systolic blood pressure and a significant increase in plasma level of apolipoprotein A. Improvement of the lipid profile in patients with cardiovascular risk have been documented in several studies [2,15,16]. In contrast to some reports in the literature, our results showed no significant




Figure 1: Effects of Ramadan fasting on glycemic control, inflammation and lipids profile markers: fasting glycaemia (mmol/L) (A), insulin (μ UI/mL) (B), HOMA-IR index (C), us-CRP(ng/L) (D), HDL-risk (E) and LDL-CT(mmol/L) (F) .

differences in HDL-C levels and apolipoprotein B levels. Contrary to our findings, Nematy et al reported a significant increase in HDL-C levels [17]. Other studies have shown that fasting is effective to ameliorate HDL, LDL/HDL, and TG/HDL ratios, and could be protective of coronary artery disease [18,19]. This contradiction may result from the differences in dietary habits and calorie intake of the study population. Different geographical locations, climate conditions, as well as different socioeconomic status in *Mediterranean populations*, may explain, in part, some discrepancies observed in these studies.

A Tunisian report showed that dietary changes in Ramadan are associated with an increase in dietary cholesterol intake, which deteriorates plasma lipids, and apolipoproteins levels in patients with atherosclerosis risk [20]. Ben Jemaa et al. showed that nutritional education programs positively impact diabetic patients who prefer RF or insist to fast during Ramadan [21].

The present study reported a decrease in us-CRP levels however; there was no significant change in homocysteine levels. This result is in contrast to that reported by Chennaoui et al, who found that CRP and homocysteine levels were significantly low during Ramadan [22]. A Tunisian pilot study in stable chronic obstructive pulmonary disease male patients identified that RF did not bring about any significant variations in CRP [23].

Although, previous investigations have indicated that RF has impressive effects on decreasing inflammation and oxidative stress markers [24,25]. Asadi and al, reported that Serum amyloid-A, a sensitive marker of an acute inflammatory state, and protein carbonyl group, an important index of oxidative stress, was decreased significantly in patients with cardiovascular disease as compared with those of the baseline (BR) [26]. Hunger

creates a stressful situation and stimulates self-regulation mechanisms with suppression of sympathetic tone [13]. This change has powerful anti-inflammatory effects which play an important role in atherogenesis and development of atherosclerotic plaques in coronary arteries [27]. Improved glycemic control has been reported in this study with a significant decrease in glycemia fasting, insulin level, and HOMA index. These results were in agreement with other studies that have demonstrated the ameliorating effect of fasting on glycemic control [28,29]. In a recent study, the majority of patients with stable ischemic heart disease and type 2 diabetes under optimal care were able to fast Ramadan without any significant change in cardiovascular risk factors or hospitalization for diabetes complications.

Glycemic control improved, however, hypoglycemia was significantly more frequent and profound compared to non- fasting state [30].

Our study has several limitations. The main one is the small number of patients included and the absence of a non-fasting control group. Insufficient data on chronobiological modifications during fasting is an important factor, which affects the association of fasting with biochemical changes and could be another limitation of the present study.

CONCLUSION

This study showed an improvement in plasma lipids and cardiometabolic risk factors in patients with stable ischemic heart disease. The majority of these patients can fast without anticipating any major adverse cardiac events.

Conflicts of Interest: The authors have no conflicts of interest to declare.

REFERENCES

1. Chamsi-Pasha M, Chamsi-Pasha H. The cardiac patient in Ramadan. *Avicenna J Med* 2016;6(2):33-38.
2. Nematy M, Alinezhad-Namaghi M, Rashed MM, Mozdehifard M, Sajjadi SS, Akhlaghi S, et al. Effects of Ramadan fasting on cardiovascular risk factors: a prospective observational study. *Nutr J* 2012;11(1). doi:10.1186/1475-2891-11-69.
3. Chamsi-Pasha H, Ahmed WH, Al-Shaibi KF. The cardiac patient during Ramadan and Hajj. *J Saudi Heart Assoc* 2014;26(4):212-15.
4. Suwaidi JA, Bener A, Hajar HA, Numan MT. Does hospitalization for congestive heart failure occur more frequently in Ramadan: a population-based study (1991-2001). *Int J Cardiol* 2004;96(2):217-21.
5. Malinowski B, Zalewska K, Weśnierska A, Sokotowska MM, Socha M, Liczner G, et al. Intermittent Fasting in Cardiovascular Disorders—An Overview. *Nutrients* 2019;11(3). doi:10.3390/nu11030673.
6. Rahman M, Rashid M, Basher S, Sultana S, Nomani MZA. Improved serum HDL cholesterol profile among Bangladeshi male students during Ramadan fasting. *East Mediterr Health J Rev Sante Mediterr Orient Al-Majallah Al-Sihhiyah Li-Sharq Al-Mutawassit* 2004;10(1-2):131-37.

7. Chennaoui M, Desgorces F, Drogou C, Boudjemaa B, Tomaszewski A, Depiesse F, et al. Effects of Ramadan fasting on physical performance and metabolic, hormonal, and inflammatory parameters in middle-distance runners. *Appl Physiol Nutr Metab Physiol Appl Nutr Metab* 2009;34(4):587-94.
8. Al Suwaidi J, Bener A, Hajar HA, Numan MT. Does hospitalization for congestive heart failure occur more frequently in Ramadan: a population-based study (1991-2001). *Int J Cardiol* 2004;96(2):217-21.
9. Campeau L. The Canadian Cardiovascular Society grading of angina pectoris revisited 30 years later. *Can J Cardiol* 2002;18:371-9.
10. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972;18(6):499-502.
11. Tang Q, Li X, Song P, Xu L. Optimal cut-off values for the homeostasis model assessment of insulin resistance (HOMA-IR) and pre-diabetes screening: Developments in research and prospects for the future. *Drug Discov Ther* 2015;9(6):380-85.
12. Chamsi-Pasha H, Ahmed WH. The effect of fasting in Ramadan on patients with heart disease. *Saudi Med J* 2004;25(1):47-51.
13. Khafaji HARH, Bener A, Osman M, Al Merri A, Al Suwaidi J. The impact of diurnal fasting during Ramadan on the lipid profile, hs-CRP, and serum leptin in stable cardiac patients. *Vasc Health Risk Manag* 2012;8:7-14.
14. Mousavi M, Mirkarimi S, Rahmani G, Hosseinzadeh E, Salehi N. Ramadan fast in patients with coronary artery disease. *Iran Red Crescent Med J* 2014;16(12):e7887.
15. Shehab A, Abdulle A, El Issa A, Al Suwaidi J, Nagelkerke N. Favorable changes in lipid profile: the effects of fasting after Ramadan. *PLoS One* 2012;7(10):e47615.
16. Unalacak M, Kara IH, Baltaci D, Erdem O, Bucaktepe PGE. Effects of Ramadan fasting on biochemical and hematological parameters and cytokines in healthy and obese individuals. *Metab Syndr Relat Disord* 2011;9(2):157-61.
17. Nematy M, Alinezhad-Namaghi M, Rashed MM, Mozhdehifard M, Sajjadi SS, Akhlaghi S, et al. Effects of Ramadan fasting on cardiovascular risk factors: a prospective observational study. *Nutr J* 2012;11:69.
18. Rahman M, Rashid M, Basher S, Sultana S, Nomani MZA. Improved serum HDL cholesterol profile among Bangladeshi male students during Ramadan fasting. *East Mediterr Health J Rev Sante Mediterr Orient Al-Majallah Al-Sihhiyah Li-Sharq Al-Mutawassit* 2004;10(1-2):131-37.
19. Akaberi A, Golshan A, Moojdekanloo M, Hashemian M. Does fasting in Ramadan ameliorate Lipid profile? A prospective observational study. *Pak J Med Sci* 2014;30(4):708-11.
20. Barkia A, Mohamed K, Smaoui M, Zouari N, Hammami M, Nasri M. Change of diet, plasma lipids, lipoproteins, and fatty acids during Ramadan: a controversial association of the considered Ramadan model with atherosclerosis risk. *J Health Popul Nutr* 2011;29(5):486-93.
21. Ben Jemaa H, Mahjoub F, Berriche O, Gammoudi A, Chaabouni S, Jamoussi H. Impacts of nutritional education before Ramadan fasting on dietary intake, weight and body composition in diabetic patients. *Tunis Med* 2019;97(10):1139-45.
22. Chennaoui M, Desgorces F, Drogou C, Boudjemaa B, Tomaszewski A, Depiesse F, et al. Effects of Ramadan fasting on physical performance and metabolic, hormonal, and inflammatory parameters in middle-distance runners. *Appl Physiol Nutr Metab Physiol Appl Nutr Metab* 2009;34(4):587-94.
23. Rejeb H, Ben Khelifa M, Ben Abdallah J, Mrad S, Ben Rejeb M, Hayouni A, et al. The Effects of Ramadan-Fasting (RF) on Inflammatory and Hematological Indices of Stable Chronic Obstructive Pulmonary Disease (COPD) Male Patients: A Pilot Study. *Am J Mens Health* 2018;12(6):2089-103.
24. Salim I, Al Suwaidi J, Ghabban W, Alkilani H, Salam AM. Impact of religious Ramadan fasting on cardiovascular disease: a systematic review of the literature. *Curr Med Res Opin* 2013;29(4):343-54.
25. Stokholm KH, Breum L, Astrup A. Cardiac contractility, central haemodynamics and blood pressure regulation during semistarvation. *Clin Physiol Oxf Engl* 1991;11(6):513-23.
26. Asadi H, Abolfathi AA, Badalzadeh R, Majidinia M, Yaghoubi A, Asadi M, et al. Effects of Ramadan Fasting on Serum Amyloid A and Protein Carbonyl Group Levels in Patients With Cardiovascular Diseases. *J Cardiovasc Thorac Res* 2015;7(2):55-59.
27. Galea J, Armstrong J, Gadsdon P, Holden H, Francis SE, Holt CM. Interleukin-1 beta in coronary arteries of patients with ischemic heart disease. *Arterioscler Thromb Vasc Biol* 1996;16(8):1000-06.
28. Khatib FA, Shafagoj YA. Metabolic alterations as a result of Ramadan fasting in non-insulin-dependent diabetes mellitus patients in relation to food intake. *Saudi Med J* 2004;25(12):1858-63.
29. Norouzy A, Mohajeri SMR, Shakeri S, Yari F, Sabery M, Philippou E, et al. Effect of Ramadan fasting on glycemic control in patients with Type 2 diabetes. *J Endocrinol Invest* 2012;35(8):766-71.
30. Hassanein M, Rashid F, Elsayed M, Basheir A, Al Saeed M, Abdelgadir E, et al. Assessment of risk of fasting during Ramadan under optimal diabetes care, in high-risk patients with diabetes and coronary heart disease through the use of FreeStyle Libre flash continuous glucose monitor (FSL-CGMS). *Diabetes Res Clin Pract* 2019;150:308-14.